

The capabilities of lidar datasets for radiative heating rate and flux calculations



Tyler Thorsen; Qiang Fu
University of Washington

David Turner

NOAA/National Severe Storms Laboratory

Rob Newsom; Jennifer Comstock

Pacific Northwest National Laboratory

Lidars and radiative fluxes


Vertical profiles of visible-wavelength extinction coefficients

Lidars and radiative fluxes

Vertical profiles of visible-wavelength extinction coefficients

- Synergy with cloud radar
 - Complementary sensitivities to cloud
(radar: poorer sensitivity; lidar: attenuation)
 - For ice, radar+lidar \rightarrow particle size
(e.g. Wang and Sassen 2002; Delanoe and Hogan 2008)
- Aerosol properties

A-train:



CALIPSO

CloudSat

ARM:



MPL


(micropulse lidar)

MMCR

(millimeter cloud radar)

- Both sets are large quantities of data, commonly used (separately)

A-train:



ARM:



CALIPSO

CloudSat


MPL

(micropulse lidar)


MMCR

(millimeter cloud radar)

- Both sets are large quantities of data, commonly used (separately)
- Compute cloud radiative effect using ARM and A-train radar+lidar observations over Darwin, Australia





Cloud radiative effect (Thorsen et al. 2013a)



- Up to 1.4 K/day difference

Cloud radiative effect (Thorsen et al. 2013a)



- Up to 1.4 K/day difference
- Mostly due to differences in lidar occurrence profiles:
MPL detects much less cirrus than CALIPSO
 - Combination of complete attenuation and a poorer sensitivity

The elastic lidar equation

We are using extinction from these lidars BUT

- Single-channel elastic backscatter lidars don't actually measure extinction

$$S_{\lambda_0}(z) \propto \beta_{p,\lambda_0}(z) \times \exp \left[- \int_0^z \alpha_{p,\lambda_0}(z') dz' \right]$$

The elastic lidar equation

We are using extinction from these lidars BUT


- Single-channel elastic backscatter lidars don't actually measure extinction

$$S_{\lambda_0}(z) \propto \beta_{p,\lambda_0}(z) \times \exp \left[- \int_0^z \alpha_{p,\lambda_0}(z') dz' \right]$$

- Assume "lidar-ratio" profile: $S_p(z) = \alpha_p(z)/\beta_p(z)$
- The lidar ratio is not constant \rightarrow large errors in extinction

The ARM Raman lidar (RL)


- Measures an elastic signal and a Raman-scattered nitrogen signal



The ARM Raman lidar (RL)


- Measures an elastic signal and a Raman-scattered nitrogen signal
- Nitrogen signal is independent of particulate backscatter

$$S_{\lambda_{N_2}}(z) \propto \exp \left[- \int_0^z \alpha_{p, \lambda_0}(z') dz' \right]$$



The ARM Raman lidar (RL)


- Measures an elastic signal and a Raman-scattered nitrogen signal
- Nitrogen signal is independent of particulate backscatter
$$S_{\lambda_{N_2}}(z) \propto \exp \left[- \int_0^z \alpha_{p, \lambda_0}(z') dz' \right]$$
- Can directly-measure cloud/aerosol extinction and backscatter



The ARM Raman lidar (RL)

- Measures an elastic signal and a Raman-scattered nitrogen signal
- Nitrogen signal is independent of particulate backscatter
$$S_{\lambda_{N_2}}(z) \propto \exp \left[- \int_0^z \alpha_{p, \lambda_0}(z') dz' \right]$$
- **Can directly-measure cloud/aerosol extinction and backscatter**
- More sensitive than the ARM MPL; cirrus occurrence agrees well with CALIPSO; cloud/aerosol detection is unbiased by the solar background

(Thorsen et al. 2013b)




The ARM Raman lidar (RL)

- Measures an elastic signal and a Raman-scattered nitrogen signal
- Nitrogen signal is independent of particulate backscatter
$$S_{\lambda_{N_2}}(z) \propto \exp \left[- \int_0^z \alpha_{p, \lambda_0}(z') dz' \right]$$
- **Can directly-measure cloud/aerosol extinction and backscatter**
- More sensitive than the ARM MPL; cirrus occurrence agrees well with CALIPSO; cloud/aerosol detection is unbiased by the solar background


(Thorsen et al. 2013b)

- **Bonus: water vapor and temperature**

(Turner et al., 2002; Newsom et al., 2010)




RL-FEX (Feature detection and EXtinction retrieval)



- New automated retrieval algorithm for the ARM RL
- Comprehensively addresses the lidar retrieval problem

(Thorsen et al. 2015; Thorsen and Fu 2015)


RL-FEX (Feature detection and EXtinction retrieval)



- New automated retrieval algorithm for the ARM RL
- Comprehensively addresses the lidar retrieval problem
- **RL-FEX makes possible statistical comparisons of CALIPSO cloud/aerosol properties to an advanced lidar**
- All previous work has focused on comparisons using case studies or small sample sizes

(Thorsen et al. 2015; Thorsen and Fu 2015)

RL-FEX (Feature detection and EXtinction retrieval)



- New automated retrieval algorithm for the ARM RL
- Comprehensively addresses the lidar retrieval problem
- RL-FEX makes possible statistical comparisons of CALIPSO cloud/aerosol properties to an advanced lidar
 - All previous work has focused on comparisons using case studies or small sample sizes
- We now have a true climatology of cloud/aerosol extinction



(Thorsen et al. 2015; Thorsen and Fu 2015)

How does using extinction from an elastic lidar (e.g. CALIPSO/MPL) affect your radiative flux calculation?

How does using extinction from an elastic lidar (e.g. CALIPSO/MPL) affect your radiative flux calculation?



- ① Calculate flux using RL-FEX best-estimate extinction
- ② Calculate flux using elastic channel-only retrieved extinction (i.e. assumed lidar ratios)
 - TOA & surface aerosol/cloud radiative effect (i.e. subtract clear-sky flux)
 - Multiyear mean values

SGP site (Oklahoma): Raman vs. elastic extinction



- Aerosols: 25–30% difference
- Clouds: $\leq 15\%$ difference, some cancellation in the net



SGP site (Oklahoma): Raman vs. elastic extinction



- Aerosols: 25–30% difference
- Clouds: $\leq 15\%$ difference, some cancellation in the net
- This is a “lidar-biased” view (only $\sim 3\text{--}4$ optical depths worth of particulates)

CALIPSO and Raman lidar aerosol occurrence

CALIPSO and Raman lidar aerosol occurrence





- Transparent profiles only (laser has fully penetrated the atmosphere)
- CALIPSO detects much less aerosol than the RL

- Is the aerosol missed by CALIPSO radiatively important?

- ① Calculation using all aerosol detected by the RL
- ② Calculation using RL data, but with aerosol randomly removed (multiple times) to force a CALIPSO-like occurrence profile ("RL degraded to CALIPSO's sensitivity")

- Is the aerosol missed by CALIPSO radiatively important?

- Calculation using all aerosol detected by the RL
- Calculation using RL data, but with aerosol randomly removed (multiple times) to force a CALIPSO-like occurrence profile ("RL degraded to CALIPSO's sensitivity")



- Significant biases of $\sim 50\text{--}75\%$

Global estimates of shortwave direct aerosol effects at TOA

Global estimates of shortwave direct aerosol effects at TOA

- Passive sensors: -5 Wm^{-2} (mostly limited to clear-sky ocean)

(Yu et al. 2006 and references therein)

Global estimates of shortwave direct aerosol effects at TOA

- Passive sensors: -5 Wm^{-2} (mostly limited to clear-sky ocean)
(Yu et al. 2006 and references therein)
- Active sensors: -0.6 to -1.9 Wm^{-2} (using CALIPSO: all-sky, land+ocean, vertically-resolved)
(Oikawa et al. 2013; Matus et al. 2015)

Global estimates of shortwave direct aerosol effects at TOA

- Passive sensors: -5 Wm^{-2} (mostly limited to clear-sky ocean)
(Yu et al. 2006 and references therein)
 - Active sensors: -0.6 to -1.9 Wm^{-2} (using CALIPSO: all-sky, land+ocean, vertically-resolved)
(Oikawa et al. 2013; Matus et al. 2015)
-

Are these estimates smaller due to CALIPSO's poor sensitivity?
(causes a $\sim 70\%$ reduction at the two ARM sites)

Summary

- CALIPSO+CloudSat better suited for heating rate calculations than MPL+MMCR due to the MPL's lack of sensitivity.
- RL-FEX: new retrieval for the ARM Raman lidar. Provides directly-retrieved cloud/aerosol extinction coefficients.
- Assumptions needed to obtain extinction from elastic lidars results in $\sim 25\%$ biases in the inferred aerosol radiative effects and $\leq 15\%$ in the cloud radiative effects.
- A significant amount of aerosol goes undetected by CALIPSO
 - This lack of aerosol reduces the inferred aerosol radiative effects significantly ($\sim 50\text{--}75\%$)

Global estimates of shortwave direct aerosol effects at TOA

- Passive sensors: -5 Wm^{-2} (mostly limited to clear-sky ocean)
(Yu et al. 2006 and references therein)
- Active sensors: -0.6 to -1.9 Wm^{-2} (using CALIPSO: all-sky, land+ocean, vertically-resolved)
(Oikawa et al. 2013; Matus et al. 2015)
 - Clear-sky ocean: -2.6 to -3.2 Wm^{-2}

Radiative transfer model details

- NASA Fu-Liou, 2 streams
- Pressure / temperature / water vapor from radiosondes; standard profiles (MLS / MLW/ TROP) fill in above
- Ozone: standard profiles
- Surface albedo = 0.2
- Clouds: extinction from RL
 - Liquid: $R_e = 8\mu m$
 - Ice: $D_{ge} = 30\mu m$
- Aerosol
 - RL extinction at 355 nm
 - SGP: d'Almedia continental model
 - TWP: d'Almedia maritime model

